Thomas Youm, MD FACS 1056 Fifth Avenue NY, NY 10028 • (212)348-3636



Rehabilitation Protocol: Meniscus Allograft Transplantation

Name:	Date:
Diagnosis:	Date of Surgery:
Phase I (Weeks 0-8)	
Weightbearing:	
Weeks 0-2: Partial We	eightbearing (up to 50%)
 Weeks 2-6: Advance to 	o WBAT with crutches (d/c crutches at 4 weeks post-op if gait normalized)
 Hinged Knee Brace: worn for 	6 weeks post-op
 Locked in full extension 	n for ambulation and sleeping - remove for hygiene (Week 1)
 Locked in full extension 	n for ambulation- remove for hygiene and sleeping (Week 2)
 Set to range from 0-90° 	for ambulation-remove for hygiene and sleeping (Weeks 3-6)
 Discontinue brace at 6 v 	weeks post-op
 Range of Motion – PROM → AA 	AROM → AROM as tolerated
Weeks 0-2: Non-weight	ntbearing 0-90°
Weeks 2-8: Full non-w	veightbearing ROM as tolerated – progress to flexion angles greater than 90'
 Therapeutic Exercises 	
 Quadriceps sets, heel sl 	ides, straight leg raises, patellar mobilizations, co-contractions (Weeks 0-2
	minal knee extensions (Weeks 2-8)
 Activities in brace for fi 	rst 6 weeks – then without brace
	h flexion > 90° during Phase I
 Avoid tibial rotation f 	or first 8 weeks to protect the meniscal allograft
Phase II (Weeks 8-12)	
Weightbearing: As tolerated	
• Range of Motion – Full active	ROM
Therapeutic Exercises	
<u>=</u>	n extension exercises, begin hamstring strengthening
o Lunges – 0-90°, Leg pre	
o Proprioception exercise	
o Begin use of the station	
Phase III (Months 3-6)	
• Weightbearing: Full weightbe	earing with normal gait nattorn
0 0	
• Range of Motion – Full/Painle	SS KUM
Therapeutic Exercises Continue with good and	d hamatring atronathoning
o Continue with quad and	
o Focus on single-leg stre	ingtii
Begin jogging/runningPlyometrics and sport-s	anagifia dvilla
7	•
	ity as tolerated (6 months post-op)
Maintenance program for stren	igth and endurance
Comments:	
Frequency: times per week	Duration: weeks
Signature:	Date: